

Acute Kidney Injury Related to Immune Checkpoint Inhibitors

Shruti Gupta MD, MPH

Director of Onconeurology, BWH/DFCI

Division of Renal Medicine, Department of Medicine

Brigham and Women's Hospital and Dana-Farber Cancer Institute

Assistant Professor of Medicine

Harvard Medical School

Shruti Gupta MD, MPH



University of Virginia School of Medicine
Medicine Residency @ MGH

Nephrology Fellowship @ BWH/MGH

Assistant Professor of Medicine @ HMS

Director, Onco-Nephrology Program

@ DFCI/BWH

- Clinical focus: Onconeurology
- Research focus: Nephrotoxicity of Cancer Therapies

Disclosures

- Research funding from BTG International, AstraZeneca, Janssen
- Research funding from NIDDK (K23DK125672 and R03DK141708)
- Consultant for Secretome, Proletariat Therapeutics, BTG, Mersana Therapeutics, Alexion, MediBeacon
- Speaker fees from Springer Inc.
- Advisory Board for Orion Pharma
- President Emeritus and Founder of American Society of Onconeurology



Objectives

- *To explain key risk factors for immune checkpoint inhibitor-associated acute kidney injury (ICI-AKI)
- *To recognize the complexities associated with diagnosing and managing ICI-AKI



Case

63 y.o. female with adenocarcinoma of lung with metastases to the brain, thoracic lymph nodes, and bone, on treatment with **pembrolizumab and pemetrexed** maintenance therapy, who presented to onconeurology for evaluation and management of AKI





Case

- Previously on carboplatin/pemetrexed/pembrolizumab, and then transitioned to pemetrexed, pembrolizumab maintenance
- SCr rose from a baseline of 0.7 mg/dl prior to starting all therapy to 1.4 mg/dl
- **Urinalysis negative** for blood, protein, white blood cells, red blood cells
- Was previously taking non-steroidal anti-inflammatory drugs regularly
- **No immune-related adverse events** while on immunotherapy
- **History of proton pump inhibitor use**

What is the next step???



The Rise of Immune Checkpoint Inhibitors (ICIs)




Patients Initiated on ICIs at BWH/DFCI and MGH




Mechanism of Action of ICIs

- Block immune inhibitory checkpoints and release “brakes” on immune system
- Release of cytolytic molecules like tumor necrosis factor- α , granzyme B, interferon- γ



ICI-AKI is one of several “Immune-related Adverse Events (irAEs)”



Postow et al., N Engl J Med, 2018

Histologic Features of ICI-AKI: Initial Case Series

Clinicopathological features of acute kidney injury associated with immune checkpoint inhibitors CrossMark

see commentary on page 474

Frank B. Cortazar¹, Kristen A. Marrone², Megan L. Troxell³, Kenneth M. Ralto⁴, Melanie P. Hoenig⁴, Julie R. Brahmer², Dung T. Le², Evan J. Lipson², Ilya G. Glezerman⁵, Jedd Wolchok⁵, Lynn D. Cornell⁶, Paul Feldman⁷, Michael B. Stokes⁸, Sarah A. Zapata⁹, F. Stephen Hodi¹⁰, Patrick A. Ott¹⁰, Michifumi Yamashita¹¹ and David E. Leaf¹²

n=13 (ATIN in 12/13)

AJKD

Case Report

Association of Acute Interstitial Nephritis With Programmed Cell Death 1 Inhibitor Therapy in Lung Cancer Patients

Anushree C. Shirali, MD,¹ Mark A. Perazella, MD,¹ and Scott Gettinger, MD²

n=6 (ATIN in 6/6)

18/19 (95%) with ATIN



Histologic Features of ICI-AKI: Beyond ATIN

SHORT REPORT

Open Access



Nephrotoxicity of immune checkpoint inhibitors beyond tubulointerstitial nephritis: single-center experience

Omar Mamlouk¹, Umut Selamet², Shana Machado¹, Maen Abdelrahim³, William F. Glass⁴, Amanda Tchakarov⁴, Lillian Gaber⁵, Amit Lahoti⁶, Biruh Workeneh⁶, Sheldon Chen⁶, Jamie Lin⁶, Noha Abdel-Wahab^{7,8}, Jean Tayar⁸, Huifang Lu⁸, Maria Suarez-Almazor⁸, Nizar Tannir⁹, Cassian Yee¹⁰, Adi Diab^{10†} and Ala Abudayyeh^{6*†}

Retrospective study of all patients treated with ICIs at MD Anderson between 2008–2018 who were biopsied for AKI ($n=16$ out of 6412 treated patients)



Histologic Features of ICI-AKI: Beyond ATIN

ATIN ($n=14$)

Pauci-immune GN ($n=3$;
only 1 had +ANCA)


AA amyloid ($n=1$)

IgAN ($n=2$)

MN ($n=1$; anti-PLA2R neg)



C3GN ($n=1$)


Secondary FSGS ($n=1$)



AKI is common in patients receiving ICIs, but how much is true ICI-AKI?

The Incidence, Causes, and Risk Factors of Acute Kidney Injury in Patients Receiving Immune Checkpoint Inhibitors

Harish Seethapathy,¹ Sophia Zhao,¹ Donald F. Chute,¹ Leyre Zubiri,² Yaa Oppong ,¹ Ian Strohhahn,¹ Frank B. Cortazar,¹ David E. Leaf ,³ Meghan J. Mooradian,² Alexandra-Chloé Villani,^{4,5} Ryan J. Sullivan,² Kerry Reynolds,² and Meghan E. Sise¹



ICI-AKI: Major Consequences, Yet Key Questions Remained Unanswered

ICI-AKI can lead to:

- Discontinuation of ICI therapy
- Irreversible loss of kidney function
- Prolonged courses of immunosuppression

Existing data largely limited to single-center case series


Limited data on safety of rechallenge



Cortazar et al., KI, 2016;
Manohar et al., NDT, 2019;
Seethapathy et al., CJASN, 2019;
Meraz-Muñoz et al., JITC, 2020;
Cortazar et al., JASN, 2020

Acute kidney injury in patients treated with immune checkpoint inhibitors

Multicenter international study
429 patients with ICI-AKI
30 sites
10 countries



Data Collection

Detailed data on **429 patients diagnosed with ICI-AKI** between **2012-2020**, from **30 sites across 10 countries**:

- Demographics and comorbidities
- Concomitant treatment with nephrotoxic medications
- Prior/concomitant extrarenal irAEs
- Laboratory and biopsy data
- Treatments for ICI-AKI
- Outcomes: renal recovery, ICI-rechallenge, survival



Definitions

Criteria for ICI-AKI

Criteria 1: Increase in SCr $\geq 100\%$ from baseline OR treatment with RRT

Criteria 2: Increase in SCr $\geq 50\%$ from baseline AND at least one of the following:

- 1) ATIN on biopsy
- 2) ICI held for at least one cycle
- 3) Treatment with corticosteroids




Definitions

Control Patients: received ICIs contemporaneously but did not develop ICI-AKI


Renal Recovery: nadir SCr ≤ 1.5 times the baseline value within 90 days following ICI-AKI

Timing of ICI-AKI



***ICI-AKI developed at a median of 16 weeks (IQR, 8-32) after ICI initiation**


Severity of ICI-AKI







- **77 patients (18%) had stage 1**
- **144 (34%) had stage 2**
- **208 (49%) had stage 3, including 33 who received RRT (8% overall)**

Extrarenal Immune-Related Adverse Events

Extrarenal irAEs in 243 (57%) patients




Can We Diagnose ICI-AKI Based on Clinical or Laboratory Features?





Pathologies on Biopsy

ATIN in 125/151 (83%) of biopsied patients





Risk Factors for ICI-AKI: Why Onconeurologists Dislike PPIs

Variable	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
Age (per 10 years)	1.17 (1.04-1.31)	1.05 (0.92-1.21)
Male sex	1.16 (0.88-1.52)	1.15 (0.86-1.53)
Combination ICPI therapy	1.42 (1.01-1.98)	1.30 (0.90-1.87)
Baseline eGFR (ml/min/1.73m ²)		
≥90 (REF)	1	1
60-89	1.54 (1.13-2.10)	1.36 (0.95-1.94)
45-59	2.48 (1.59-3.87)	2.23 (1.35-3.68)
<45	1.92 (1.74-4.89)	2.62 (1.47-4.65)
PPI use*	2.55 (1.92-3.40)	2.40 (1.79-3.23)
Prior or concomitant extrarenal irAEs**	2.19 (1.65-2.91)	2.07 (1.53-2.78)



Treatment with Corticosteroids




350 (82%) patients were treated with steroids, of whom 100 received IV pulse dose steroids



Recovery and Rechallenge

- Renal recovery in 276 patients (64.3%)
- Early treatment with corticosteroids is associated with a higher odds of renal recovery
- Of 121 patients rechallenged, fewer than 1 in 5 patients had recurrent ICI-AKI after rechallenge



Back to Our Case

Previously on carboplatin/pemetrexed/pembrolizumab, and then transitioned to pemetrexed, pembrolizumab maintenance
SCr rose from a baseline of 0.7 mg/dl prior to starting all therapy to 1.4 mg/dl

Urinalysis negative for blood, protein, WBCs, RBCs

Was previously taking NSAIDs regularly

No immune-related adverse events while on immunotherapy

History of proton pump inhibitor use

What is the next step???





Biopsy:

- *Hematuria, proteinuria (particularly albuminuria), +serologies
- ***Those receiving combination therapy**
- *ANY concern for alternative cause




No biopsy


- *Solitary kidney/history of nephrectomy
- *Anticoagulation
- *No obvious alternative cause
- *Frailty; patient preference

Novel Biomarkers of ICI-AKI


Urine Biomarkers




Retinol-binding Protein/Cr*CRP



Serum Biomarkers




C Reactive Protein (CRP)



Moledina DG et al. J Clin Invest. 2023.
Sise ME et al. J Immunother Cancer. 2023.

Farooqui et al. KI Reports. 2022
Isik B et al. KI Reports. 2021.
Courtesy of Abhijat Kitchlu

Novel Biomarkers of ICI-AKI: Urinary CXCL-9



Beyond Urinary and Blood-Based Biomarkers...


F¹⁸-FDG PET imaging as a diagnostic tool for immune checkpoint inhibitor-associated acute kidney injury

Shruti Gupta,^{1,2,3} Olivia Green-Lingren,¹ Sudhir Bhimaniya,^{3,4} Aleksandra Krokmal,⁴ Heather Jacene,^{3,4} Marlies Ostermann,⁵ Sugama Chicklore,⁶ Ben Sprangers,^{7,8} Christophe M. Deroose,⁹ Sandra M. Herrmann,¹⁰ Sophia L. Wells,¹ Sarah A. Kaunfer,¹ Jessica L. Ortega,¹ Clara García-Carro,¹¹ Michael Bold,¹² Kevin L. Chen,¹³ Meghan E. Sise,^{3,14} Pedram Heidari,¹⁵ Wai Lun Will Pak,¹⁶ Meghan D. Lee,¹⁴ Pazit Beckerman,¹⁷ Yael Eshet,¹⁸ Raymond K. Hsu,¹⁹ Miguel Hernandez Pampaloni,²⁰ Arash Rashidi,²¹ Norbert Avril,²² Vicki Donley,²¹ Zain Mithani,²³ Russ Kuker,²⁴ Muhammad O Awiwi,²⁵ Mindy X. Wang,²⁶ Sujal I. Shah,²⁷ Michael D. Weintraub,^{3,28} Heiko Schoder,²⁹ Raad B. Chowdhury,^{1,2,3} Harish Seethapathy,^{3,14} Kerry L. Reynolds,^{3,30} Maria Jose Soler,³¹ Ala Abudayyeh,²⁶ Ilya Glezerman,¹⁶ and David E. Leaf^{1,3}

Published August 8, 2024 - [More info](#)




FDG-PET/CTs: Percentage change in SUV_{mean} from baseline to the time of AKI



- ICI-AKI: the SUV_{mean} increased by a median of **57.4% (IQR, 40.3 to 119.8)** from baseline to follow-up
- AKI from non-ICI causes: SUV_{mean} increased by 8.5% (IQR, 1.4 to 19.9)
- Controls without AKI: SUV_{mean} decreased by 0.8% (IQR, -16.6 to 5.1)

AUC for Differentiation of ICI-AKI from Controls




AUC 0.97 (95% CI, 0.93-1.00)

AUC unchanged after a sensitivity analysis where we excluded patients in the control group if they were receiving AIN-causing medications



Representative Image of a Baseline and Follow-Up PET-CT



Case Follow-Up

63 y.o. female with adenocarcinoma of lung with metastases to the brain, thoracic lymph nodes, and bone, on treatment with **pembrolizumab and pemetrexed** maintenance therapy, who presented to onconeurology for evaluation and management of AKI



Case Follow-Up

Previously on carboplatin/pemetrexed/pembrolizumab, and then transitioned to pemetrexed, pembrolizumab maintenance
SCr rose from a baseline of 0.7 mg/dl prior to starting all therapy to 1.4 mg/dl

Urinalysis negative for blood, protein, WBCs, RBCs

Was previously taking NSAIDs regularly


No immune-related adverse events while on immunotherapy

History of proton pump inhibitor use



Case Follow-Up

Decision was made to pursue a kidney biopsy....





Case Follow-Up

- Started on prednisone 1 mg/kg while awaiting biopsy
- Slow improvement in kidney function with return of serum creatinine to 0.95 mg/dl
- Successfully rechallenged, OFF proton pump inhibitor, and without recurrence of ICI-AKI



Treatment of ICI-AKI with Steroids: How Long is Enough?



Special Populations

- Kidney Transplant
- Underlying Autoimmune Diseases



A multi-center study on safety and efficacy of immune checkpoint inhibitors in cancer patients with kidney transplant.

Retrospective cohort study (2010-2020)



International
Multi-center
(23 institutions)



Kidney transplant
recipients
(n=69)



ICI therapy for
advanced cancer
(aPD-1, aPD-L1,
aCTLA-4)

Safety



Acute rejection
42%



Time to rejection
24 days



Graft loss
65% of rejection

Efficacy: Tumor response to ICI therapy (complete response + partial response)











Skin squamous cell carcinoma (n=24)
36%

Melanoma (n=22)
40%

CONCLUSION:

Immune checkpoint inhibitors are associated with high acute rejection rate but result in reasonable tumor response.

⑧ Cemiplimab for Kidney Transplant Recipients With Advanced Cutaneous Squamous Cell Carcinoma

Glenn J. Hanna, MD¹ ; Harita Dharanesswaran, BS² ; Anita Giobbie-Hurder, MS³ ; John J. Harran, RN²; Zixi Liao, RN²; Lori Pai, MD⁴; Vatche Tchekmedyan, MD⁵ ; Emily S. Ruiz, MD² ; Abigail H. Waldman, MD²; Chrysalyne D. Schmults, MD² ; Leonardo V. Riella, MD, PhD⁶ ; Patrick Lizotte, PhD⁷ ; Cloud P. Paweletz, PhD⁷; Anil K. Chandraker, MD, MBCHB⁸; Naoka Murakami, MD, PhD⁸ ; and Ann W. Silk, MD² 

- Phase 1 study of cemiplimab for kidney transplant recipients with metastatic SCC (n=12)
- After cross-taper to an mTOR inhibitor and pulsed dose corticosteroids, patients received cemiplimab intravenously once every 3 weeks for up to 2 years and were assessed for response every 8 weeks.
- Primary end point was the rate of kidney rejection
- No kidney rejection or loss was observed
- Response was observed in 5 of 11 evaluable patients, including 2 with durable responses

Autoimmune Disease

- ICI use is feasible in patients with pre-existing autoimmune disorders who develop cancer, though disease activity, level of immunosuppression, and the extent of organ damage and function should be considered
- AIM-NIVO trial underway




ICI-AKI Position Statement

Expert panel consensus
recommendations for diagnosis and
management of ICI-AKI



Herrmann SM , Abudayyeh A, Gupta S, et al.
Kidney Int. 2024.



Question #1: Which of the Following is not Known to Be a Risk Factor for ICI-AKI?

- A. Proton pump inhibitor use
- B. Allopurinol use
- C. Lower Baseline eGFR
- D. Extrarenal immune-related adverse events



Question #1: Which of the Following is not Known to Be a Risk Factor for ICI-AKI?

- A. Proton pump inhibitor use
- B. Allopurinol use
- C. Lower Baseline eGFR
- D. Extrarenal immune-related adverse events



Question #2: Which of the Following is the Most Common Pathology on Kidney Biopsy in a Patient with ICI-AKI?

- A. Acute tubular injury
- B. IgA nephropathy
- C. Acute tubulointerstitial nephritis
- D. Thrombotic microangiopathy



Question #2: Which of the Following is the Most Common Pathology on Kidney Biopsy in a Patient with ICI-AKI?

- A. Acute tubular injury
- B. IgA nephropathy
- C. Acute tubulointerstitial nephritis
- D. Thrombotic microangiopathy



Question #3: A 65 y/o patient with metastatic melanoma is referred to you for the evaluation of AKI. He has been treated with nivolumab for the past 4 months. He has not had any known exposure to proton pump inhibitors, and he has no history of any extrarenal irAEs. His SCr has risen from a prior baseline of 1.0 to 1.5 mg/dl. He is found to have 2+ blood and 2+ protein on urinalysis. A 24-hour urine protein collection reveals 2.4 g of proteinuria, of which 2 g is albuminuria. Renal US shows normal-sized kidneys and no evidence of obstruction. On examination of his urine sediment, he has 10-12 non-dysmorphic RBCs per hpf.



What is your next step in the management of this patient?

- A. Empiric treatment with corticosteroids
- B. Empiric treatment with rituximab
- C. Perform a kidney biopsy
- D. Continue immunotherapy since the AKI is not severe
- E. Check a soluble IL2 receptor, and if elevated, start corticosteroids



What is your next step in the management of this patient?

- A. Empiric treatment with corticosteroids
- B. Empiric treatment with rituximab
- C. Perform a kidney biopsy**
- D. Continue immunotherapy since the AKI is not severe
- E. Check a soluble IL2 receptor, and if elevated, start corticosteroids



Acknowledgements

- Co-PI of ICI-AKI consortium, Dr. David E. Leaf
- Collaborators at all of our sites around the world
- My small but mighty lab (shrutiGkidney.org)
- Leaders and founding members of the American Society of Onconeurology
- My husband and my 2 daughters (oldest wants to be a kidney doctor!)



References

- 1) Gupta S, Short SA, Sise M. Acute kidney injury in patients treated with immune checkpoint inhibitors. *J Immunother Cancer*. 2021;9:e003467.
- 2) Sise ME, Wang Q, Seethapathy H, Moreno D, Harden D, Smith RN, Rosales IA, Colvin RB, Chute S, Cornell LD, Herrmann S, Fadden R, Sullivan RJ, Yang N, Barmettler S, Wells SL, Gupta S, Villani AC, Reynolds K, Farmer J. Soluble and Cell-Based Markers of Immune Checkpoint Inhibitor Associated Nephritis. *J Immunother Cancer*. 2023; 11(1): e006222.
- 3) Cortazar FB, Marrone KA, Troxell ML. Clinicopathological features of acute kidney injury and checkpoint inhibitors. *Kidney Int* 206;90:638-47.
- 4) Shirali AC, Perazella MA, Gettinger S. Association of acute interstitial nephritis with programmed cell death 1 inhibitor therapy in lung cancer patients. *Am J Kidney Dis* 2016;68:287-91.
- 5) Isik B, Alexander MP, Manohar S. Biomarkers, clinical features, and rechallenge for immune checkpoint inhibitor renal immune-related adverse events. *Kidney Int Rep* 202;6:1022-31.
- 6) Mamlouk O, Selamet U, Machado S, et al. Nephrotoxicity of immune checkpoint inhibitors beyond tubulointerstitial nephritis: a single-center experience. *J Immunotherapy Cancer* 2019;7.
- 7) Seethapathy J, Zhao S, Chute DF. The incidence, causes, and risk factors for acute kidney injury in patients receiving immune checkpoint inhibitors. *Clin Journ Amer Soc Neph*. 2019; 14(12): 692-700.
- 8) Herrmann SM, Abudayyeh A, Gupta S, et al. Diagnosis and Management of Immune Checkpoint Inhibitor-Associated Nephrotoxicity: A Position Statement from the American Society of Onco-nephrology. *Kidney Int*. 2024; 107(1):21-32.
- 9) Gupta S,* Green-Lingren O,* Bhimaniya S,* et al. F18-FDG PET imaging as a diagnostic tool for immune checkpoint inhibitor-associated acute kidney injury. *J Clin Invest*. 2024; 134(18).
- 10) Gupta S, Garcia-Carro C, Prosek JM, et al. Shorter versus Longer Corticosteroid Duration and Recurrent Immune Checkpoint Inhibitor-Associated AKI. *J Immunother Cancer*. 2022; 10(9):e005646.
- 11) Gupta S,* Strohbehn IA,* Wang Q, et al. Acute Kidney Injury in Patients Receiving Pembrolizumab Combination Therapy versus Pembrolizumab Monotherapy for Advanced Lung Cancer. *Kidney Int*. 2022;102(4): 930-935.

